

Dosing Information for Healthcare Professionals

Concentrated
MOTRIN®
Infants' Drops

Children's
MOTRIN®

Use this chart to determine the proper dose of MOTRIN® for your patient.

If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.
Do not give more than 4 doses in 24 hours.**

**Concentrated MOTRIN®
Infants' Drops**

Active Ingredient:
Ibuprofen 50 mg (NSAID)*
(in each 1.25 mL)

**Children's MOTRIN®
Oral Suspension**

Active Ingredient:
Ibuprofen 100 mg (NSAID)*
(in each 5 mL or 1 tsp)

DO NOT GIVE if under 6 mos old

DO NOT GIVE if under 6 mos old

1.25 mL

—

1.875 mL

—

—

5 mL (1 tsp)

—

7.5 mL (1½ tsp)

—

10 mL (2 tsp)

—

12.5 mL (2½ tsp)

—

15 mL (3 tsp)

WEIGHT	AGE
—	<6 mos
12-17 lbs	6-11 mos
18-23 lbs	12-23 mos
24-35 lbs	2-3 yrs
36-47 lbs	4-5 yrs
48-59 lbs	6-8 yrs
60-71 lbs	9-10 yrs
72-95 lbs	11 yrs

Remind parents and caregivers:

- Read and follow the label on all MOTRIN® products
- Repeat dose every 6-8 hours as needed
- Do NOT give more than 4 doses in 24 hours
- Use only the dosing device (syringe or dosing cup) that came with the product

*Nonsteroidal anti-inflammatory drug

mL = milliliter

tsp = teaspoon