

# Dosing Information for Healthcare Professionals

Infants'  
**TYLENOL**

Children's  
**TYLENOL**

Use this chart to determine the proper dose of TYLENOL<sup>®</sup> for your patient.

**If possible, use weight to dose; otherwise use age.**

WEIGHT		AGE	
6-11 lbs	2-3 mos	<b>1.25 mL</b>	—
12-17 lbs	4-11 mos	<b>2.5 mL</b>	—
18-23 lbs	12-23 mos	<b>3.75 mL</b>	—
24-35 lbs	2-3 yrs	<b>5 mL</b>	<b>5 mL</b> (1 tsp)
36-47 lbs	4-5 yrs	—	<b>7.5 mL</b> (1½ tsp)
48-59 lbs	6-8 yrs	—	<b>10 mL</b> (2 tsp)
60-71 lbs	9-10 yrs	—	<b>12.5 mL</b> (2½ tsp)
72-95 lbs	11 yrs	—	<b>15 mL</b> (3 tsp)

**DOSE: Every 4 hours as needed.  
Do not give more than 5 doses in 24 hours.**

### Infants' TYLENOL<sup>®</sup> Oral Suspension

**Active Ingredient:**  
Acetaminophen  
160 mg  
(in each 5 mL)

### Children's TYLENOL<sup>®</sup> Oral Suspension

**Active Ingredient:**  
Acetaminophen  
160 mg  
(in each 5 mL or 1 tsp)

### Remind parents and caregivers:

- Read and follow the label on all TYLENOL<sup>®</sup> products
- Repeat dose every 4 hours while symptoms last
- Do NOT give more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device (syringe or dosing cup) that came with the product

**All Infants' TYLENOL<sup>®</sup> and Children's TYLENOL<sup>®</sup> Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)**

mL = milliliter

tsp = teaspoon